

# Nodal Solutions Bifurcating from Infinity for some Singular $p$ -Laplace Equations: Flat and Compact Support Solutions

Jesùs I. Díaz

*Instituto de Matemática Interdisciplinar,  
Universidad Complutense de Madrid, 28040 Madrid, Spain  
ildefonso\_diaz@mat.ucm.es*

Jesùs Hernández

*Don Ramón de la Cruz 73-7<sup>o</sup> A, 28001 Madrid, Spain*

Francisco J. Mancebo

*E.T.S. Ingeniería Aeronáutica y del Espacio,  
Universidad Politécnica de Madrid, 28040 Madrid, Spain  
fj.mancebo@upm.es*

Received: July 23, 2015

Accepted: November 9, 2015

We study the existence and multiplicity of nodal solutions with normal exterior derivative different or equal to zero (case of *flat solutions*) or having a *free boundary* (the boundary of the set where the solution vanishes) of some one-dimensional  $p$ -Laplace problems of eigenvalue type with homogeneous Dirichlet boundary conditions and a, possibly singular, nonlinear absorption term.

*2000 Mathematics Subject Classification:* 35J65, 35B32, 35P30, 35R35.

## 1. Introduction

This paper deals with the study of the countable branches of nodal solutions bifurcating from the infinity for the one-dimensional nonlinear eigenvalue problem

$$\begin{cases} -\frac{d}{dx} \left( \left| \frac{du}{dx} \right|^{p-2} \frac{du}{dx} \right) + |u|^{m-1}u = \lambda |u|^{p-2}u & \text{in } ]-1, 1[, \\ u(-1) = u(1) = 0, \end{cases} \quad (1)$$

where  $p > 1$ ,  $\lambda$  is a positive parameter,  $m$  and  $p$  are given real numbers such that

$$-1 < m < p - 1. \quad (2)$$

We point out that the above differential equation must be slightly modified for some values of the parameter  $m$ . So, for  $m = 0$  the equation should be understood in the framework of the multivalued maximal monotone graphs of  $\mathbb{R}^2$  as

$$-(|u'|^{p-2} u')' + H(u) \ni \lambda |u|^{p-2}u$$

where  $H(u)$  is the multivalued maximal monotone graph of  $\mathbb{R}^2$  given by

$$H(r) = \begin{cases} -1 & \text{if } r < 0, \\ [-1, 1] & \text{if } r = 0, \\ +1 & \text{if } r > 0. \end{cases}$$

This case (at least for  $p = 2$ ) is specially relevant in image processing (see [18], [19]). Moreover, if  $m \in (-1, 0)$  then the equation should be replaced by

$$-(|u'|^{p-2} u')' + |u|^{m-1} u \chi_{\{u \neq 0\}} = \lambda |u|^{p-2} u$$

where

$$\chi_{\{u \neq 0\}}(x) = \begin{cases} 1 & \text{if } u(x) \neq 0, \\ 0 & \text{if } u(x) = 0, \end{cases}$$

and thus the whole expression  $|u|^{m-1} u \chi_{\{u \neq 0\}}$  must be understood as

$$[|u|^{m-1} u \chi_{\{u \neq 0\}}](x) = \begin{cases} |u(x)|^{m-1} u(x) & \text{if } u(x) \neq 0, \\ 0 & \text{if } u(x) = 0. \end{cases}$$

We study this problem by using phase plane methods for ordinary differential equations. This kind of arguments were used by the authors in [7] and [11]. These methods provide a complete description of the solution set for (1).

The related problem for the equation

$$\begin{cases} -\frac{d}{dx} \left( \left| \frac{du}{dx} \right|^{p-2} \frac{du}{dx} \right) + |u|^{m-1} u = \lambda |u|^{q-1} u & \text{in } ]-1, 1[, \\ u(-1) = u(1) = 0, \end{cases} \quad (3)$$

with  $0 < m < q < p-1$  was studied initially in [7], for  $p = 2$ , giving a complete description of both positive solutions and the continua with infinitely many compact support solutions obtained from the very especial solution satisfying condition (7) below by some “stretching” manipulations. These results were extended to the  $p$ -Laplacian allowing singular nonlinearities satisfying  $-1 < m < q < p-1$  in [11] by using similar (more sophisticated) arguments. Then the case of a general bounded domain  $\Omega \subset \mathbb{R}^N$  and  $p = 2$  was treated by Il'yasov and Egorov in [17] combining variational and continuation methods, by using this time a variant of the Mountain Pass Theorem. The study of the stability of solutions was made in [10]. For related work see [16], [24] and [25].

The above problem (1) was also considered for  $p = 2$  and  $0 < m < 1$  in [8] again with ODE methods giving sharper results than the ones obtained through abstract bifurcation tools by Rabinowitz [22], [21], [23]. We point out that although there are several results dealing with the bifurcation from the infinity for the  $p$ -Laplace operator (see, e.g. Drabek et al. [12]) our results are new in this direction.

Besides many related problems in the literature (see, e.g., references in the monographs [4], [1]) a relevant motivation for this work was to provide some kind of

“alternative” approach to the “ambiguity” raised by some compactly supported solutions to the linear Schrödinger equation on the real line (see [5], [6] and [8] for more details and references). There is also an extensive literature on the Schrödinger quasilinear equation associated to the  $p$ -Laplace diffusion operator (see e.g. You et al. [26], and its many references). The association of problems as (1) with the study of standing waves for the related semilinear Schrödinger equation and the semilinear wave equation with real coefficients was also largely considered in the literature (at least for  $p = 2$ : see, e.g. [2] and its references). Nevertheless, in both types of equations (Schrödinger and wave equation) the consideration of the case  $p \neq 2$  leads to some new facts that seem not to be well presented in the previous literature (see Remark 2.8 below).

As mentioned before, similar results giving a complete description of non-negative (including positive) solutions together with the transition to compact support solutions were given in [8] for the case  $p = 2$ . For the case of a bounded domain  $\Omega \subset \mathbb{R}^N$ , a partial result (namely the existence of an unbounded continuum of non-negative solutions bifurcating from infinity at  $\lambda_1 > 0$ , the first eigenvalue to  $-\Delta$  with Dirichlet boundary conditions) was obtained in [14] using asymptotic bifurcation arguments. Much later, Porretta proved in [20] the existence of (at least) a non-negative solution for any  $\lambda > \lambda_1$  (for  $p = 2$ ) and this result was improved in [9] using this time variational methods (Nehari manifolds) and a Pohozaev identity.

The contents of this paper is the following: in Section 2 we analyse the existence of a branch of nonnegative solutions for a bounded interval of the parameter  $\lambda \in ]\lambda_1(p), \lambda_1^*(m, p)[$ . We recall that  $\lambda_1(p)$  is the first eigenvalue of the nonlinear equation

$$\begin{cases} -\frac{d}{dx} \left( \left| \frac{du}{dx} \right|^{p-2} \frac{du}{dx} \right) = \lambda |u|^{p-2} u & \text{in } ]-1, 1[ , \\ u(-1) = u(1) = 0. \end{cases} \tag{4}$$

The first eigenvalue of the  $p$ -Laplacian for one dimensional domains was obtained by Otani in [13] and its value is given by

$$\lambda_1(p) = (p - 1) \left( \frac{\pi}{p \sin\left(\frac{\pi}{p}\right)} \right)^p. \tag{5}$$

Here  $\lambda_1^*(m, p)$  is a certain value of the parameter whose exact definition depends crucially on  $p$  and  $m$ . The value of  $\lambda_1^*(m, p)$  can be written in closed form as

$$\lambda_1^*(m, p) = (p - 1) \left( \frac{\pi}{(p - m - 1) \sin\left(\frac{\pi}{p}\right)} \right)^p = \left( \frac{p}{p - m - 1} \right)^p \lambda_1(p). \tag{6}$$

We show that the unique positive solution of (1) for  $\lambda = \lambda_1^*(m, p)$  has a particular behaviour at  $x = \pm 1$ . Due to this behaviour this solution is called as “flat solu-

tion” (and also as “free boundary nonnegative solution” by other authors) since although  $u(x) > 0$  for any  $x \in (-1, 1)$  it satisfies

$$\frac{\partial u}{\partial n}(\pm 1) = 0. \quad (7)$$

We also give some estimates on the convergence

$$\frac{\partial u_\lambda}{\partial n}(\pm 1) \nearrow 0 \quad \text{as} \quad \lambda \searrow \lambda_1^*(m, p),$$

and make explicit some non-degeneracy estimates for  $u_{\lambda_1^*(m, p)}$  (near the boundary of the interval  $\Omega = (-1, 1)$ ) extending the estimates given in ([6] for  $p = 2$ ). The associated solution  $u_{\lambda_1^*(m, p)}$  (when extended by zero to the real line  $\mathbb{R}$ ) gives rise to a continuum of nonnegative solutions  $u_\lambda$  for  $\lambda > \lambda_1^*(m, p)$  through an appropriate rescaling. This kind of solutions have compact support included in  $[-1, 1]$ .

In section 3 we obtain a qualitatively similar result for the branches of nodal solutions with a finite number of simple zeroes and that bifurcate from infinity. Here we will denote by  $\lambda_k$  the bifurcation value of the branch that bifurcates from the infinity and possesses  $(k - 1)$  sign-changes. Now the values of  $\lambda_k$  and  $\lambda_k^*(m, p)$  can be written in closed form as

$$\lambda_k(p) = (p - 1) \left( \frac{k\pi}{p \sin\left(\frac{\pi}{p}\right)} \right)^p = k^p \lambda_1(p), \quad (8)$$

and

$$\begin{aligned} \lambda_k^*(m, p) &= (p - 1) \left( \frac{k\pi}{(p - m - 1) \sin\left(\frac{\pi}{p}\right)} \right)^p \\ &= \left( \frac{p}{p - m - 1} \right)^p \lambda_k(p) = k^p \lambda_1^*(m, p). \end{aligned} \quad (9)$$

We show that the behavior of the nodal solutions near the points where they vanish is of the same type that the behaviour of nonnegative solutions near the boundary.

## 2. The branch of nonnegative solutions.

In this Section we study the non-negative solutions of the equation (1) by using ordinary differential equations arguments. This allows us to obtain a complete description of the solution set as a function of the parameter  $\lambda$ . We generalize the results obtained by Díaz and Hernández in [8] for the case  $p = 2$  and  $m > 0$ . Now we consider the equation (1) under the assumption (2).

We recall that, as in [11], once we define

$$f(u) = \lambda|u|^{p-2}u - |u|^{m-1}u\chi_{\{u \neq 0\}},$$

(if  $m \neq 0$ ) we can introduce the following notions of solution:

**Definition 2.1.** We say that  $u \in W_0^{1,p}(-1,1)$  is a *positive strong solution* of problem (1) if  $u > 0$  on  $(-1,1)$ ,  $f(u) \in L^1(-1,1)$ ,  $(|u'|^{p-2}u')' \in L^1(-1,1)$ ,  $-(|u'|^{p-2}u')'(x) = f(u(x))$  for a.e.  $x \in (-1, 1)$  and  $\frac{\partial u}{\partial n}(\pm 1) < 0$ .

**Definition 2.2.** We say  $u \in W_0^{1,p}(-1,1)$  is a *flat solution* of problem (1) if  $u$  is as in the preceding definition but replacing the last condition by  $\frac{\partial u}{\partial n}(\pm 1) = 0$ .

In the case  $m = 0$  the above definitions must be adapted in the sense that now it must exist  $h \in L^1(-1,1)$ , with  $h(x) \in H(u(x))$  for a.e.  $x \in (-1, 1)$  such that  $-(|u'|^{p-2}u')'(x) + h(x) = \lambda|u(x)|^{p-2}u(x)$  for a.e.  $x \in ]-1, 1[$ .

The main result of this section is the following:

**Theorem 2.3.** *Let  $\lambda_1(p)$  and  $\lambda_1^*(m, p)$  be given by (5) and (6) respectively. Then:*

- a) *If  $\lambda \in ]0, \lambda_1(p)[$  there is no nonnegative solution of (1).*
- b) *If  $\lambda \in ]\lambda_1(p), \lambda_1^*(m, p)[$  there is a unique positive solution  $u_\lambda$  of (1). Moreover,*

$$\frac{\partial u_\lambda}{\partial n}(\pm 1) < 0 \tag{10}$$

and 
$$\underline{K}d(x, \partial\Omega) \leq u_\lambda(x) \leq \overline{K}d(x, \partial\Omega), \tag{11}$$

$$\Omega = (-1, 1), \text{ for some constants } \overline{K} > \underline{K} > 0.$$

- c) *If  $\lambda = \lambda_1^*(m, p)$  there is a unique positive solution  $u_{\lambda_1^*(m,p)}$  of (1). Moreover,*

$$\frac{\partial u_{\lambda_1^*(m,p)}}{\partial n}(\pm 1) = 0, \tag{12}$$

$$\|u_{\lambda_1^*(m,p)}\|_{L^\infty[-1,1]} = \left( \frac{p}{(m+1)\lambda_1^*} \right)^{\frac{1}{p-m-1}},$$

and 
$$\underline{K}d(x, \partial\Omega)^{p/(p-1-m)} \leq u_{\lambda_1^*(m,p)}(x) \leq \overline{K}d(x, \partial\Omega)^{p/(p-1-m)} \tag{13}$$

$$\Omega = (-1, 1), \text{ for some constants } \overline{K} > \underline{K} > 0.$$

- d) *If  $\lambda > \lambda_1^*(m, p)$  then the function  $u_{\lambda,\zeta} : [-1, 1] \rightarrow \mathbb{R}$  defined by*

$$u_{\lambda,\zeta}(x) = \begin{cases} \left( \frac{\lambda_1^*(m,p)}{\lambda} \right)^{\frac{1}{p-m-1}} u_{\lambda_1^*(m,p)}\left(\frac{x-\zeta}{\omega}\right) & \text{if } |x-\zeta| < \omega, \\ 0 & \text{if } x \in [-1, \zeta-\omega] \cup [\zeta+\omega, 1], \end{cases}$$

*is a nonnegative solution of (1), where  $\omega = \left( \frac{\lambda_1^*(m,p)}{\lambda} \right)^{\frac{1}{p}} < 1$ ,  $\zeta \in [-1 + \omega, 1 - \omega]$  and  $u_{\lambda_1^*(m,p)}$  is the unique positive solution of (1) for  $\lambda = \lambda_1^*(m, p)$ . Consequently, for each  $\lambda > \lambda_1^*(m, p)$  there is a family of nonnegative solutions of (1),  $u_{\lambda,\zeta}$ , that depends arbitrarily on the parameter  $\zeta \in [-1 + \omega, 1 - \omega]$ . Moreover, the behavior of  $u_{\lambda,\zeta}$  near the boundary of the points where  $u_{\lambda,\zeta}$  vanishes is given by (13).*

For the proof we shall need the following auxiliary Lemma.

**Lemma 2.4.** *Let  $\gamma : [r_F, +\infty[ \rightarrow \mathbb{R}$  be the function defined by*

$$\gamma(\mu) = \int_0^\mu \sqrt[p]{\frac{p-1}{p}} \frac{1}{\sqrt[p]{F(\mu) - F(r)}} dr,$$

where  $r_F = \left(\frac{p}{m+1}\right)^{\frac{1}{p-m-1}}$ , and  $F(r) = \left(\frac{r^p}{p} - \frac{r^{m+1}}{m+1}\right)$ . The function  $\gamma$  is such that

i)  $\gamma \in C([r_F, +\infty[) \cap C^1(]r_F, +\infty[)$ .

ii)  $\frac{d\gamma}{d\mu}(\mu) < 0$  for all  $\mu \in ]r_F, +\infty[$ .

iii)  $\lim_{\mu \rightarrow r_F^+} \frac{d\gamma}{d\mu}(\mu) = -\infty$  if  $\frac{p}{p+1} \leq m+1$  and  $\frac{d\gamma}{d\mu}(r_F^+)$  is finite if  $m+1 < \frac{p}{p+1}$ .

iv)  $\gamma(r_F) = \frac{\pi \sqrt[p]{p-1}}{(p-m-1) \sin\left(\frac{\pi}{p}\right)}$ ,  $\lim_{\mu \rightarrow +\infty} \gamma(\mu) = \sqrt[p]{p-1} \left(\frac{\pi}{p \sin\left(\frac{\pi}{p}\right)}\right)$ .

**Remark 2.5.** A curious fact is that in [15] (respectively [11]) it was shown the increasing nature of the curve  $\gamma$  (which also implies the uniqueness of solution) for problem (3) when  $m$  is near  $-1$ ,  $p = 2$  (respectively  $p \neq 2$ ) and  $q \in (m, 1)$  (respectively  $q \in (m, p-1)$ ).

**Proof of Lemma 2.4.** We use the change of variables  $\tau := r/\mu$  in the definition of  $\gamma$ . Then

$$\begin{aligned} \gamma(\mu) &= \sqrt[p]{p-1} \int_0^\mu \frac{1}{\sqrt[p]{\mu^p - \frac{p}{m+1}\mu^{m+1} - \left(r^p - \frac{p}{m+1}r^{m+1}\right)}} dr \\ &= \sqrt[p]{p-1} \int_0^1 \frac{1}{\sqrt[p]{1 - \tau^p - \frac{p}{(m+1)\mu^{p-m-1}}(1 - \tau^{m+1})}} d\tau. \end{aligned} \quad (14)$$

We have

$$\frac{d\gamma}{d\mu}(\mu) = -\sqrt[p]{p-1} \int_0^1 \frac{\frac{p-m-1}{(m+1)\mu^{p-m}}(1 - \tau^{m+1})}{\sqrt[p]{\left(1 - \tau^p - \frac{p}{(m+1)\mu^{p-m-1}}(1 - \tau^{m+1})\right)^{p+1}}} d\tau. \quad (15)$$

For  $\mu \in ]r_F, \infty[$  it is not difficult to verify that the integral in (15) is convergent,

and thus  $\frac{d\gamma}{d\mu}(\mu) \in C(r_F, \infty)$  with  $\frac{d\gamma}{d\mu}(\mu) < 0$ . According to the assumption (2)

the integral (15) is divergent for  $\mu = r_F$ , hence  $\lim_{\mu \rightarrow r_F^+} \frac{d\gamma}{d\mu}(\mu) = -\infty$ .

On the other hand, from assumption (2), the integral (14) is bounded as  $\mu \downarrow r_F$ , and in conclusion  $\gamma \in C([r_F, \infty))$ . By virtue of the Fatou's lemma:

$$\begin{aligned} \lim_{\mu \rightarrow +\infty} \gamma(\mu) &= \lim_{\mu \rightarrow +\infty} \sqrt[p]{p-1} \int_0^1 \frac{1}{\sqrt[p]{1-\tau^p - \frac{p}{(m+1)\mu^{p-m-1}}(1-\tau^{m+1})}} d\tau \\ &= \sqrt[p]{p-1} \int_0^1 \frac{1}{\sqrt[p]{1-\tau^p}} d\tau = \frac{\sqrt[p]{p-1}}{p} \int_0^1 \sigma^{\frac{1}{p}-1} (1-\sigma)^{1-\frac{1}{p}-1} d\sigma \\ &= \frac{\sqrt[p]{p-1}}{p} B\left(\frac{1}{p}, 1-\frac{1}{p}\right). \end{aligned}$$

Taking into account that  $B(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$  and  $\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)}$  (see e.g. [3]) we obtain

$$\lim_{\mu \rightarrow +\infty} \gamma(\mu) = \frac{\sqrt[p]{p-1}}{p} \frac{\pi}{\sin\left(\frac{\pi}{p}\right)}.$$

Finally

$$\begin{aligned} \gamma(r_F) &= \sqrt[p]{p-1} \int_0^1 \frac{1}{\sqrt[p]{\tau^{m+1} - \tau^p}} d\tau = \sqrt[p]{p-1} \int_0^1 \frac{1}{\tau^{\frac{m+1}{p}} \sqrt[p]{1-\tau^{p-m-1}}} d\tau \\ &= \frac{\sqrt[p]{p-1}}{p-m-1} \int_0^1 \sigma^{\frac{1}{p}-1} (1-\sigma)^{1-\frac{1}{p}-1} d\sigma = \frac{\sqrt[p]{p-1}}{p-m-1} B\left(\frac{1}{p}, 1-\frac{1}{p}\right) \\ &= \frac{\sqrt[p]{p-1}}{(p-m-1)} \frac{\pi}{\sin\left(\frac{\pi}{p}\right)}. \quad \square \end{aligned}$$

**Remark 2.6.** Note that for  $p = 2$  and  $m = 0$  the function  $\gamma : [2, +\infty[ \rightarrow \mathbb{R}$  can be written in closed form as

$$\gamma(\mu) = \frac{\pi}{2} + \arcsin\left(\frac{1}{\mu-1}\right).$$

**Proof of Theorem 2.3.** Upon multiplication of equation (1) by  $u$ , integrating by parts and taking into account the variational definition of the first eigenvalue of (4) we get part a).

To show the qualitative behavior of solutions of the equation (1), we make the change of variables

$$y = Lx, \quad u\left(\frac{y}{L}\right) = \left(\frac{1}{L}\right)^{\frac{p}{p-m-1}} w(y), \quad \text{with } \lambda = L^p. \quad (16)$$

in equation (1) to obtain

$$\mathcal{P}(L) \begin{cases} -\frac{d}{dy} \left( \left| \frac{dw}{dy} \right|^{p-2} \frac{dw}{dy} \right) + |w|^{m-1}w = |w|^{p-2}w \quad \text{in } ]-L, L[, \\ w(-L) = w(L) = 0. \end{cases} \quad (17)$$

If a positive solution of  $\mathcal{P}(L)$  exists then necessarily it will have a maximum  $\mu > 0$  at some point  $\zeta \in (-L, L)$ . So, let us consider

$$\mathcal{CP} \begin{cases} -(|u'|^{p-2} u')' = -|u|^{m-1}u + |u|^{p-2}u \\ u(\zeta) = \mu, \quad u'(\zeta) = 0. \end{cases}$$

Hence, we have to consider only the case  $\mu \in [r_F, \infty)$ . Notice that, if  $u$  is a strong positive solution of  $\mathcal{P}(L)$  then  $(|u'|^{p-2} u')' \in L^1(-L, L)$  implies that  $u' \in L^\infty(-L, L)$ . Then, since by definition  $f \in L^1(-L, L)$ , the formula

$$\begin{aligned} -\int_{\zeta}^x (|u'|^{p-2} u')' u' d\tau &= -\int_{\zeta}^x \frac{p-1}{p} \frac{d}{dy} (|u'|)^p(\tau) d\tau \\ &= \int_{\zeta}^x f(u(\tau)) u'(\tau) d\tau = \int_{\zeta}^x F'(u(\tau)) d\tau, \end{aligned}$$

is well justified. Since  $F'(s) = f(s) = -|s|^{m-1}s + |s|^{p-2}s > 0$  if  $s > 1$  and  $F(s) = \frac{s^p}{p} - \frac{s^{m+1}}{m+1} > 0$  if  $s \in ]r_F, +\infty[$  we have that for any  $x \in (-L, L)$

$$|u'(x)| = \sqrt[p]{\frac{p}{p-1} (F(\mu) - F(u(x)))}. \quad (18)$$

Notice that, in fact  $u'(x) \leq 0$  for  $x \in (-L, L)$  near  $\zeta$  with  $x > \zeta$  and that  $u'(x) \geq 0$  for  $x$  near  $\zeta$  and  $x < \zeta$ .

The solution of this equation is implicitly defined by

$$\int_{u(x)}^{\mu} \frac{dr}{\sqrt[p]{\frac{p}{p-1} (F(\mu) - F(r))}} = |\zeta - x|, \quad (19)$$

since the singularity at  $r = \mu$  is integrable, i.e.,

$$\int_s^{\mu} \frac{dr}{(F(\mu) - F(r))^{1/p}} < \infty \quad (20)$$

for any  $s \in (\mu - \varepsilon, \mu)$  and for any small enough  $\varepsilon > 0$ .

Indeed, it is easy to check that (20) always holds since  $F(\mu) - F(r) \geq \delta(\mu - r)$  for some  $\delta > 0$  and for any  $r$  near  $\mu$ . Moreover, for a strong positive solution  $u$  of problem  $\mathcal{CP}$ ,  $u = 0$  only at  $r = \pm L$ .

To finish the proof we only need to justify that the function  $u(x)$  defined implicitly by (19) satisfies that  $f(u) \in L^1(-L, L)$ . But we know that for  $x \in (-L, -L + \delta)$ ,  $u(x) \in [0, \varepsilon)$  for some  $\varepsilon > 0$ , small enough, and we get that

$$-C \int_{-L}^{-L+\delta} f(u(x))dx \leq - \int_{-L}^{-L+\delta} f(u(x))u'(x)dx = \int_{-L}^{-L+\delta} |u'|^p dx < \infty$$

for some  $C$  near  $\sqrt[p]{\frac{p}{p-1}F(\mu)}$ , since  $u \in C^1(-L, L)$ , and the proof of the existence of part b) ends.

As a byproduct we obtain that  $u$  is an even function. Equation (19), when particularized for  $x = -L$  and  $x = L$ , gives the identity  $|\zeta + L| = |\zeta - L|$ , which implies  $\zeta = 0$ , particularizing again (19) for  $x$  and  $-x$  we obtain  $u(x) = u(-x)$ , i. e.  $u$  is an even function.

In the case of  $\lambda = \lambda^*(m)$  the associated function  $v$  is such that  $\mu = r_F$  and in consequence  $v'(\pm L) = 0$ . Moreover, since

$$\frac{1}{m+1}s^{1+m} \geq \frac{s^{m+1}}{m+1} - \frac{s^p}{p} \geq \frac{(p-1-m)}{p(1+m)}s^{1+m} \quad \text{for } s \in (0, 1),$$

we get that there exist two positive constants  $\underline{M} < \overline{M}$  such that

$$\underline{M}\tau^{\frac{p-1-m}{p}} \leq \frac{1}{\sqrt[p]{\frac{p}{p-1}}} \int_0^\tau \frac{dr}{\sqrt[p]{-F(r)}} \leq \overline{M}\tau^{\frac{p-1-m}{p}} \tag{21}$$

for any  $\tau \in (0, 1)$  which leads to conclusion (13).

The dependence on  $\lambda$  of the value of the normal derivative of  $u_\lambda$  at the boundary can be obtained from (18):

$$\left| \frac{du_\lambda}{dx}(\pm 1) \right| = \left( \frac{1}{\lambda} \right)^{\frac{m+1}{p(p-m-1)}} \sqrt[p]{\frac{p}{p-1}F(\gamma^{-1}(\lambda^{\frac{1}{p}}))}. \tag{22}$$

Since the function  $F$  is increasing in  $[1, +\infty[$ ,  $\gamma^{-1}([\gamma(+\infty), \gamma(r_F)]) = [r_F, +\infty[ \subset [1, +\infty]$ , and the functions  $\gamma^{-1}$  and  $\left(\frac{1}{s}\right)^{\frac{m+1}{p(p-m-1)}}$  are decreasing,  $\left| \frac{du_\lambda}{dx}(\pm 1) \right|$  is a decreasing function of  $\lambda$ . In addition  $F(\gamma^{-1}((\lambda_1^*(m, p))^{\frac{1}{p}})) = 0$  which implies that  $\frac{du_\lambda}{dx}(\pm 1) = 0$  if and only if  $\lambda = \lambda_1^*(m, p)$ . (Notice that  $F \circ \gamma^{-1}$  is differentiable only if  $m + 1 < \frac{p}{p+1}$ ).

The proof of part d) is similar to the the proof of part v) of Theorem 1 in [11].

According to (19) the unique positive solution of  $\mathcal{P}(L)$  satisfies the identity  $\gamma(w(0)) = L$ , where  $\gamma$  is as defined in Lemma 1. Since  $\gamma$  is injective and continuous, its inverse  $\gamma^{-1} : ] \lim_{\mu \rightarrow +\infty} \gamma(\mu), \gamma(r_F) ] \rightarrow \mathbb{R}$  is well defined, and gives the bifurcation diagram of the equation  $\mathcal{P}(L)$ . Hence the bifurcation diagram of (1), when written in terms of  $\|u\|_{L^\infty([-1,1])}$  and  $\lambda$ , is given by

$$\|u\|_{L^\infty([-1,1])} = \frac{\gamma^{-1}(\lambda^{\frac{1}{p}})}{\lambda^{\frac{1}{p-m-1}}}. \quad \square$$

**Remark 2.7.** For  $p = 2$  and  $m = 0$  the bifurcation diagram of (1) can be written in closed form as

$$\|u\|_{L^\infty([-1,1])} = \frac{1}{\lambda} \left( 1 + \frac{1}{\sin(\sqrt{\lambda} - \frac{\pi}{2})} \right),$$

where  $\lambda \in ]\lambda_1(2), \lambda_1^*(0, 2) ] \equiv ]\frac{\pi^2}{4}, \pi^2]$ .

**Remark 2.8.** For both types of wave equations (the Schrödinger and the semi-linear wave equation) mentioned in the Introduction the consideration of the case  $p \neq 2$  leads to some new facts that seem to be not well mentioned in the previous literature. Indeed: if we consider, for instance,  $m \in (0, 1)$ , the study of standing wave solutions leads to the consideration of the problem  $-(|u'|^{p-2} u')' + |u|^{m-1} u = \lambda u$ , and thus, by the results of [11], the difference with respect to the case  $p = 2$  is that there is an additional strictly positive solution (that does not exist for  $p = 2$ ) which is stable (since this part of the bifurcation diagram of solutions is increasing in  $\lambda$ ).

### 3. The branches of nodal solutions

In this section we consider solutions of (1) that change sign. Let us define precisely what we mean by nodal solutions.

**Definition 3.1.** Let  $Z$  be the set  $\{x_1, \dots, x_k : -1 = x_0 < x_1 < x_2 < \dots < x_{k-1} < x_k < x_{k+1} = 1\}$ . A solution of (1),  $u_\lambda : [-1, 1] \rightarrow \mathbb{R}$  is a nodal solution with  $k$  nodes if  $u_\lambda$  satisfies the conditions of Definition 1 except the positivity condition,  $u_\lambda(x) = 0$  if  $x \in Z$  and  $u_\lambda(x) \neq 0$  if  $x \in ]-1, 1[$  and  $x \notin Z$ .

The main result of this section is the following:

**Theorem 3.2.** Let  $\lambda_k(p)$  and  $\lambda_k^*(m, p)$  be the positive real numbers defined by (8) – (9) with  $k \geq 2$ . Then:

- a) If  $\lambda \in ]0, \lambda_k(p) [ \equiv ]0, k^p \lambda_1(p) [$  there is no nodal solution of (1) with  $(k-1)$  nodes.
- b) If  $\lambda \in ]\lambda_k(p), \lambda_k^*(m, p) [ \equiv ]k^p \lambda_1(p), k^p \lambda_1^*(m, p) [$  there is a unique nodal solution (1) with  $(k-1)$  nodes. Moreover

$$\frac{\partial u_\lambda}{\partial n}(x_i) \neq 0, \quad \text{for } i \in \{0, 1, \dots, k+1\}$$

and 
$$\underline{K}|x - x_i| \leq |u_\lambda(x)| \leq \overline{K}|x - x_i|, \tag{23}$$

if  $|x - x_i| = d(x, Z)$ , for some constants  $\overline{K} > \underline{K} > 0$ .

c) If  $\lambda = \lambda_k^*(m, p) \equiv k^p \lambda_1^*(m, p)$  there is only one nodal solution,  $u_{\lambda_k^*(m, p)}$  (1) with  $(k - 1)$  nodes. Moreover

$$\frac{\partial u_{\lambda_k^*(m, p)}}{\partial n}(x_i) = 0, \quad \text{for } i \in \{0, 1, \dots, k + 1\}.$$

and

$$\|u_{\lambda_k^*(m, p)}\|_{L^\infty[-1, 1]} = \left( \frac{p}{(m + 1)\lambda_k^*} \right)^{\frac{1}{p-m-1}}.$$

Moreover, the behavior of  $u_{\lambda_k^*(m, p)}$  near the boundary of the points where  $u_{\lambda_k^*(m, p)}$  vanishes is of the same type than (13), i.e.

$$\underline{K}|x - x_i|^{p/(p-1-m)} \leq |u_{\lambda_k^*(m, p)}(x)| \leq \overline{K}|x - x_i|^{p/(p-1-m)} \tag{24}$$

if  $|x - x_i| = d(x, Z)$ , for some constants  $\overline{K} > \underline{K} > 0$ .

d) If  $\lambda > \lambda_k^*(m, p)$  then the function  $u_\lambda : [-1, 1] \rightarrow \mathbb{R}$  defined by

$$u_\lambda(x) = \sum_{i=1}^{i=k} s(i)u_{\lambda, \zeta_i}(x)$$

is a solution of (1) where

$$u_{\lambda, \zeta_i}(x) = \begin{cases} \left( \frac{\lambda_1^*(m, p)}{\lambda} \right)^{\frac{1}{p-m-1}} u_{\lambda_1^*(m, p)}\left(\frac{x - \zeta_i}{\omega}\right) & \text{if } |x - \zeta_i| < \omega, \\ 0 & \text{if } x \in [-1, \zeta_i - \omega] \cup [\zeta_i + \omega, 1], \end{cases}$$

$$\omega = \left( \frac{\lambda_1^*(m, p)}{\lambda} \right)^{\frac{1}{p}}, \quad \zeta_i \in [-1 + \omega, 1 - \omega] \text{ with } \zeta_i \in [-1 + \omega, 1 - \omega] \text{ for } i = 1, \dots, k$$

and  $\zeta_{i+1} - \zeta_i > 2\omega$  for  $i = 1, \dots, k - 1$ ,  $u_{\lambda_1^*(m, p)}$  is the unique positive solution of (1) for  $\lambda = \lambda_1^*(m, p)$ , and consider the set of functions  $s : \{1, 2, \dots, k\} \rightarrow \{-1, 1\}$  such that  $s(1) = 1$  and that exists  $i \in \{2, \dots, k\}$  such that  $s(i) = -1$ . Therefore, for each  $\lambda > \lambda_k^*(m, p)$  there are, up to a change sign,  $2^{(k-1)} - 1$  continua of nonnegative-nonpositive solutions that depend arbitrarily on the parameters  $\zeta_i$  with  $i \in \{1, \dots, k\}$ . The number of functions  $s$  defined above, up to a change sign, is  $2^{(k-1)} - 1$ , where  $k > 1$ . Moreover the behavior of  $u_\lambda$  near the boundary of the points where  $u_\lambda$  vanishes is given by (24).

The following two lemmata show that the structure of these branches can be obtained from the branch of positive solutions.

**Lemma 3.3.** *Let  $u_\mu : [-1, 1] \rightarrow \mathbb{R}$  be a nodal solution of (1) for  $\lambda = \mu$  with  $k$  nodes. The function  $u_{\mu,i}^+ : [-\frac{x_i - x_{i-1}}{2}, \frac{x_i - x_{i-1}}{2}] \rightarrow \mathbb{R}$  defined by  $u_{\mu,i}^+(x) = |u_\mu(x - \frac{x_i + x_{i-1}}{2})|$  is a positive solution of*

$$-\frac{d}{dx} \left( \left| \frac{du}{dx} \right|^{p-2} \frac{du}{dx} \right) + |u|^{m-1} u = \mu |u|^{p-2} u, \text{ in } ] -\frac{x_i - x_{i-1}}{2}, \frac{x_i - x_{i-1}}{2} [, \quad (25)$$

$$u(-\frac{x_i - x_{i-1}}{2}) = u(\frac{x_i - x_{i-1}}{2}) = 0,$$

where  $i \in \{1, \dots, k, k + 1\}$ . Moreover, the zeros of  $u_\lambda$  are equispaced in  $[-1, 1]$ , i.e.,  $x_i - x_{i-1} = \frac{2}{k + 1}$  and the function  $w_\lambda : [-1, 1] \rightarrow \mathbb{R}$  defined by

$$w_\lambda(y) = (k + 1)^{\frac{p}{p-m-1}} u_{\mu,i}^+(y \frac{x_i - x_{i-1}}{2}) \quad \text{for } y \in [-1, 1],$$

is the unique positive solution of (1) for  $\lambda = \frac{\mu}{(k + 1)^p}$ .

**Proof of Lemma 3.3.** Since the differential equation (1) is invariant under translations, under the transformation  $(u) \rightarrow (-u)$  and  $u_\mu$  satisfies  $u_\mu(x_{i-1}) = u_\mu(x_i) = 0$ , the function  $u_{\mu,i}^+$  is a solution of (25) for all  $i \in \{1, \dots, k, k + 1\}$ . In addition,  $u_\mu$  is continuous and by (18)  $u \in C^1(]-1, 1[)$ , consequently

$$\frac{du_{\mu,i}^+}{dx}(x_i^-) = -\frac{du_{\mu,i+1}^+}{dx}(x_i^+).$$

As we have proved in Lemma 2.4,  $u_{\mu,i}^+$ , for all  $i \in \{1, \dots, k, k + 1\}$ , are even functions and according to (18) the maxima of each of them are equal. Therefore, the uniqueness for positive solutions implies the identity  $x_i - x_{i-1} = x_{i+1} - x_i$ , i.e, the zeroes of a nodal solution divide the interval  $]-1, 1[$  in  $k + 1$  subintervals of the same length, in other words  $x_i - x_{i-1} = 2/(k + 1)$  where  $i \in 1, \dots, k$  and  $x_0 = -1$ . Finally, if we make the change of independent variable

$$y = \frac{2x}{x_i - x_{i-1}} \equiv (k + 1)x, \quad w_\lambda(y) = (k + 1)^{\frac{p}{p-m-1}} u_{\mu,i}^+ \left( \frac{y}{k + 1} \right)$$

in (25) we obtain that  $w_\lambda$  is a solution of (1) for  $\lambda = \frac{\mu}{(k + 1)^p}$ . □

**Lemma 3.4.** *Let  $u_\lambda : [-1, 1] \rightarrow \mathbb{R}$  be a nodal solution of (1) with  $k$  nodes. If  $\frac{du_\lambda}{dx}(x_i) \neq 0$  then  $u_\lambda$  is such that*

$$u_\lambda(x + \frac{2i}{k + 1}) = -u_\lambda(-x + \frac{2i}{k + 1}) \quad (26)$$

for all  $x \in [0, \frac{2}{k + 1}]$  and all  $i \in \{1, \dots, k\}$ .

**Proof of Lemma 3.4.** According to Theorem 2.3 the function  $u_\lambda$  belongs to  $C^1(\cdot - 1, \cdot + 1)$ ,  $u_{\lambda,i}^+$  is a solution of (25) and symmetric with respect to  $(x_i + x_{i-1})/2$ . Since the positive solution is unique for (25), is invariant under translations, for the transformations  $u \rightarrow -u$  and  $x \rightarrow -x$  the identity (26) follows.  $\square$

**Proof of Theorem 3.2.** Combining the results in Theorem 2.3, Lemma 3.3 and Lemma 3.4, Theorem 3.2 follows.  $\square$

### Acknowledgements

The research of J.I. Díaz and J. Hernández was partially supported by the projects MTM2011-26119 and MTM2014-57113 of the DGISPI (Spain). The research of J.I. Díaz was partially supported by the UCM Research Group MOMAT (Ref. 910480).

### References

- [1] S. N. Antontsev, J. I. Díaz, S. I. Shmarev: *Energy methods for free boundary problems: Applications to nonlinear PDEs and fluid mechanics*, Birkhäuser-Verlag, Basel (2001).
- [2] M. Balabane, J. Dolbeault, H. Ounaies: *Nodal solutions for a sublinear elliptic equation*, *Nonlinear Anal. TMA* 52 (2003) 219–237.
- [3] P. J. Davis: *Gamma function and related functions*, in: M. Abramowitz and I. A. Stegun (eds.), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover, New York (1972) 253–294.
- [4] J. I. Díaz: *Nonlinear partial differential equations and free boundaries*, Pitman, London (1985).
- [5] J. I. Díaz: *Lectures on the Schrödinger equation for the “infinite well potential”*, in: Book of abstracts of the congresses on Quasilinear PDE’s, Tours (France), June 4-6 (2011); Partial Differential Equations, Valencia (Spain) July 1-5 (2013); Free Boundary Programme of the Newton Institute, Cambridge (UK), May 22 (2014).
- [6] J. I. Díaz: *On the ambiguous treatment of the Schrödinger equation for the infinite potential well and an alternative via flat solutions: the one-dimensional case*, *Interfaces and Free Boundaries* 17(3) (2015) 333–351.
- [7] J. I. Díaz, J. Hernández: *Global bifurcation and continua of nonnegative solutions for a quasilinear elliptic problem*, *C. R. Acad. Sci. Paris, Serie I* 329 (1999) 587–592.
- [8] J. I. Díaz, J. Hernández: *On branches of nodal solutions bifurcating from infinity for a semilinear equation related to the linear Schrödinger equation: free boundary solutions and solutions with compact support*, *Portugaliae Mathematica* 72(2-3) (2015) 145–160.
- [9] J. I. Díaz, J. Hernández, Y. Il’yasov: *On the existence of positive solutions and solutions with compact support for a spectral nonlinear elliptic problem with strong absorption*, *Nonlinear Anal.* 119 (2015) 484–500.

- [10] J. I. Díaz, J. Hernández, Y. Il'yasov: *Flat solutions of some non-Lipschitz autonomous semilinear equations may be stable for  $N \geq 3$* , submitted.
- [11] J. I. Díaz, J. Hernández, F. J. Mancebo: *Branches of positive and free boundary solutions for some singular quasilinear elliptic problems*, *J. Math. Anal. Appl.* 352 (2009) 449–474.
- [12] P. Drábek, P. Girg, P. Takac, M. Ulm: *The Fredholm alternative for the  $p$ -Laplacian: bifurcation from infinity, existence and multiplicity of solutions*, *Indiana Univ. Math. J.* 53(2) (2004) 433–482.
- [13] M. Ôtani: *A remark on certain nonlinear elliptic equations*, *Proceedings of the Faculty of Science, Tokai University*, 19 (1984) 23–28.
- [14] J. Hernández: *Bifurcación y soluciones positivas para algunos problemas de tipo unilateral*, Ph. D. Thesis, Universidad Autónoma de Madrid (1977).
- [15] T. Horváth, P. L. Simon: *On the exact number of solutions of a singular boundary value problem*, *J. Diff. Int. Equations* 22 (2009) 787–796.
- [16] P. Huang, S. H. Wang, T. Yeh: *Classification of bifurcation diagrams of a  $p$ -Laplacian semipositone problem*, *Communications on Pure and Appl. Analysis* 12(5) (2013) 2297–2318.
- [17] Y. Il'yasov, Y. Egorov: *Hopf boundary maximum principle violation for semilinear elliptic equations*, *Nonlinear Anal. TMA* 72(7-8) (2010) 3346–3355.
- [18] V. Ozolins, R. Lai, R. Caffisch, S. Osher: *Compressed modes for variational problems in mathematics and physics*, *Proc. Natl. Acad. Sci. USA* 110(46) (2013) 18368–18373.
- [19] V. Ozolins, R. Lai, R. Caffisch, S. Osher: *Compressed plane waves yield a compactly supported multiresolution basis for the Laplace operator*, *Proc. Natl. Acad. Sci. USA* 111(5) (2014) 1691–1696.
- [20] A. Porretta: *A note on the bifurcation of solutions for an elliptic sublinear problem*, *Rend. Sem. Mat. Univ. Padova* 107 (2002) 153–164.
- [21] P. H. Rabinowitz: *A global theorem for nonlinear eigenvalue problems and applications*, in: E. H. Zaratonello (ed.), *Contributions to Nonlinear Functional Analysis*, Academic Press, New York (1971) 11–36.
- [22] P. H. Rabinowitz: *Some global results for nonlinear eigenvalue problems*, *J. Funct. Anal.* 7 (1971) 487–513.
- [23] P. H. Rabinowitz: *On bifurcation from infinity*, *J. Differential Equations* 14 (1973) 462–475.
- [24] S. H. Wang, T. S. Yeh: *A complete classification of bifurcation diagrams of a Dirichlet problem with concave-convex nonlinearities*, *J. Math. Anal. Appl.* 291 (2004) 128–153.
- [25] S. H. Wang, T. S. Yeh: *Classification of bifurcation diagrams of a  $p$ -Laplacian Dirichlet problem with examples*, *J. Math. Anal. Appl.* 369 (2010) 188–204.
- [26] B. You, Y. Hou, F. Li, J. Jiang: *Pullback attractors for the non-autonomous quasi-linear complex Ginzburg-Landau equation with  $p$ -Laplacian*, *Discrete Cont. Dynam. Systems Ser. B* 19(6) (2014) 1801–1814.